



Composition and functional diversity of zooplankton-associated Prokaryote communities

Dziallas, Claudia; Grossart, Hans-Peter; Nielsen, Torkel Gissel; Tang, Kam W.

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Dziallas, C., Grossart, H-P., Nielsen, T. G., & Tang, K. W. (2012). *Composition and functional diversity of zooplankton-associated Prokaryote communities*. Poster session presented at 2012 ASLO Aquatic Sciences Meeting, Salt Lake City, UT, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Composition and Functional Diversity of Zooplankton-associated Prokaryote Communities in a Greenlandic Fjord

Claudia Dziallas¹, Hans-Peter Grossart^{1,2}, Torkel G. Nielsen^{3,4}, Kam W. Tang^{5,*}

¹Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Neuglobsow, Germany

²Potsdam University, Institute for Biochemistry and Biology, Potsdam, Germany

³National Institute of Aquatic Resources, DTU-Aqua, Technical University of Denmark, Denmark

⁴Greenland Climate Research Centre, Greenland Institute of Natural Resources, Nuuk, Greenland


⁵Virginia Institute of Marine Science, College of William & Mary, Virginia (*kamtang@vims.edu)

Background

- Prokaryotes densely colonize zooplankton bodies (Tang 2005, Grossart et al. 2009), but they are often poorly sampled or characterized (Tang et al. 2010).

- Greenlandic fjords are very productive ecosystems, but information on free-living and zooplankton-associated prokaryotes there is very limited.

- We compared the prokaryote communities in pelagic water samples and those associated with zooplankton along a salinity gradient in Godthåbsfjord, Greenland (Tang et al. 2011a).




Methods

- Surface water and zooplankton samples were collected during the ECOGREEN (2008) and BOFYGO (2010) expeditions in the spring on board R/V Dana.

- Prokaryote community compositions were analyzed by DGGE and clone libraries.

- PCR screening was done to look for specific functional groups.

Multi-dimensional scaling analysis of prokaryote community compositions




- Calanus*, *Metridia* (calanoid copepods) and water samples formed distinct clusters in terms of prokaryote compositions.

- Calanus* feeds mainly on diatoms in the spring; *Metridia* feeds omnivorously throughout the year.

- Dietary differences could lead to delivery of different prokaryotes to the two copepod species (Tang 2005, Tang et al. 2009).

Phylogenetic tree of prokaryotes in zooplankton and water samples



PCR screening for specific prokaryote functional groups

Sample type		<i>Calanus hyperboreus</i>	<i>Pareuchaeta sp.</i>	<i>Metridia longa</i>	<i>Chaetognath</i>	Water
Prokaryote group	Functional gene					
Methanogens	<i>mcrA</i>	✓		(✓)	✓	
Sulfate reducers	<i>dsr</i>	✓			✓	
Nitrogen fixers	<i>nifH</i>	✓			✓	
Dentrifiers	<i>nirK</i>				✓	
Denitrifiers	<i>nirS</i>				✓	
NH ₃ -oxidizing archaea	<i>amoA</i>	✓				✓
Planctomycetes		✓	✓		✓	
Archaea		✓	✓			✓

- Some prokaryote groups were found exclusively in zooplankton.

- Functional genes indicated occurrence of anaerobic microbial processes in zooplankton: methanogenesis, sulfate reduction, nitrogen fixation and denitrification.

Statistical analysis of environmental factors structuring prokaryotic community composition in the water column

	all	salinity	temperature	fluorescence	density	depth	station
Bacteria	R=0.246 p<0.001	R=0.12 p=0.014	R=0.011 p=0.388	R=0.134 p=0.008	R=0.113 p=0.018	R=0.116 p=0.011	R=0.52 p<0.001
Archaea	R=0.137 p=0.018	R=0.143 p=0.020	R=0.051 p=0.203	R=-0.01 p=0.528	R=0.13 p=0.022	R=0.115 p=0.006	R=0.116 p=0.025
α-Proteobacteria	R=0.211 p<0.001	R=0.282 p<0.001	R=0.113 p=0.029	R=0.087 p=0.056	R=0.276 p<0.001	R=0.175 p<0.001	R=0.427 p<0.001
Actinobacteria	R=0.073 p=0.088	R=0.047 p=0.173	R=0.028 p=0.284	R=-0.018 p=0.616	R=0.042 p=0.185	R=0.043 p=0.139	R=0.262 p<0.001

- Mantel-Test revealed spatial variation of community structure of Bacteria, α-Proteobacteria, and Archaea along the salinity gradient.

- Density, depth and station also influenced prokaryotic community structure.

Summary

- Zooplankton-associated prokaryotes increased the overall prokaryote diversity in the water column, consistent with studies in other aquatic systems (Grossart et al. 2009, Tang et al. 2009).


- Anoxic zooplankton guts may support anaerobic microbial activities within the oxygenated water column (Tang et al. 2011b), which was supported by our PCR functional gene analysis.

- Potential changes in plankton compositions in Greenland fjords in response to rapid climate change could lead to corresponding changes in prokaryote diversity and biogeochemical functions.

Outreach Efforts

- Plankton and bacteria are not familiar to Greenlanders, but they are important for sustaining Greenland fisheries.

- We worked with reporters to disseminate our research via internet blogs and local newspaper articles.



- Microbes are not “charismatic megafauna” that K-12 students normally encounter in their curricula.
- Tang worked with his graduate student Samantha Bickel through the NSF-funded **GK-12 PERFECT** project at VIMS to bring microbial ecology into grade 10 classrooms.



GK-12 Fellow in action
Graduate student Samantha Bickel teaches high school students about her research on marine microbes.
For more information, please visit:
<http://www.vims.edu/education/teaching/gk12/>

Related Oral Presentation

Zooplankton-associated bacterial abundance and community composition in the York River tributary of Chesapeake Bay
Samantha L. Bickel, Kam W. Tang, Hans-Peter Grossart [Abstract 11633; Tuesday 9:15 am, session 153 Ballroom D]

References and Acknowledgements

Grossart HP, Dziallas C, Tang KW (2009) *Environmental Microbiology Reports* 1:50-55
Tang KW (2005) *Aquatic Microbial Ecology* 38:31-40
Tang K, Dziallas C, Hutalle-Schmelzer K, Grossart H-P (2009) *Biology Letters* 5:549-553
Tang KW, Nielsen TG, Munk P, Mortensen J, Møller EF, Arendt KE, Tønnesson K, Pedersen TJ (2011a) *Marine Ecology Progress Series* 434:77-90
Tang KW, Glud R, Glud A, Rysgaard S, Nielsen TG (2011b) *Limnology and Oceanography* 56:666-672
Tang KW, Turk V, Grossart H-P (2010) *Aquatic Microbial Ecology* 61:261-277
Funding was provided by: Commission for Scientific Research in Greenland, the Danish Natural Sciences Research Council (ECOGREEN and BOFYGO); Greenland Climate Research Center; Humboldt Research Fellowship for Experienced Researchers; U.S. National Science Foundation (OCE-0814558 and DGE-0840804).